

NO 0155 002


THE CHEMISTRY OF PRECURSORS TO STEROID CARCINOGENS
DIEGO STATE UNIV CA DEPT OF CHEMISTRY M A RING ET AL
25 MAR 88 AFOSR-TR-88-0520 AFOSR-83-0209

UNCLASSIFIED

F/G 7/2

ML





REPORT DOCUMENTATION PAGE

CTIONS
TING FORM
NUMBER

2

AD-A193 062

1. REPORT NUMBER

AFOSR-TR-88-0520

4. TITLE (and Subtitle)

The Chemistry of Precursors to Silicon Carbide

5. TYPE OF REPORT & PERIOD COVERED

Final Report

6. PERFORMING ORG. REPORT NUMBER

7. AUTHOR(s)

M. A. Ring and H. E. O'Neal

8. CONTRACT OR GRANT NUMBER(s)

AFOSR 83-0209

9. PERFORMING ORGANIZATION NAME AND ADDRESS

Department of Chemistry
San Diego State University
San Diego, CA 9218210. PROGRAM ELEMENT, PROJECT, TASK
AREA & WORK UNIT NUMBERS

61102F 2303/B2

11. CONTROLLING OFFICE NAME AND ADDRESS

Air Force Office of Scientific Research
Building 410
Bolling AFB, DC 20332

12. REPORT DATE

3/25/88

13. NUMBER OF PAGES

14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)

AFOSR/NC
Bolling AFB DC 20332-6448 (Bldg 410)

15. SECURITY CLASS. (of this report)

UNCLASSIFIED

16. DECLASSIFICATION/DOWNGRADING
SCHEDULE

16. DISTRIBUTION STATEMENT (of this Report)

Approved for Public Release

Distribution Unlimited

DTIC

ELECTE

MAY 04 1988

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20 (if different from Report))

18. SUPPLEMENTARY NOTES

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Silane, Organosubstituted Mono and Disilanes, Silylene, Organosubstituted
Silylenes

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

The research can be divided into 4 main areas:

- 1) Mechanisms and Kinetics of the Decompositions of Silane and Organosubstituted Monosilanes, and of Disilane and Organosubstituted Disilanes.
- 2) Relative Rate Kinetics of the Trapping Reactions of SiH_2 with Ethylene, Acetylene, Butadiene, Methanol, Silane and 1-butene.

Cont. (See Reverse Side)

DD FORM 1 JAN 73 1473

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. ABSTRACT (Cont.)

- 3) Mechanisms and Kinetics of the Isomerizations and Decompositions of Organosilylenes.
- 4) From the above studies, we were able to propose a general reaction mechanism for carbosilane production from the decomposition of dimethylsilane (paper #12).

FINAL PROJECT REPORT (AFOSR-83-0209)

1. SUMMARY OF WORK

Our AFOSR research has focused on three important areas of organosilicon chemistry as that chemistry relates to the formation and production of carbosilanes:






Area #1) Mechanisms and Kinetics of the Decompositions of Silane and Organosubstituted Monosilanes, and of Disilane and Organosubstituted Disilanes.

Systems investigated were n-propylsilane¹, ethylsilane², silane^{3,4}, dimethylsilane^{5,12}, ethynylsilane⁶, butylsilane⁷, chloromethyldimethylsilane⁸, and 1,1,2,2-tetramethyldisilane¹³, 1,1,2-trimethyldisilane¹⁴ and 1,1-dimethyldisilane¹⁵.

Results of area #1 studies: The main findings of these studies were:

- 1) The decomposition kinetics of monosubstituted silanes (RSiH_3), are independent of the nature of the R group, (A-factors $\approx 10^{15.2 \pm 0.2} \text{ sec}^{-1}$ and $E_a \approx 64 \pm 1 \text{ Kcal}$).
- 2) The primary dissociation reaction channel of RSiH_3 compounds is 1,1 H_2 elimination (rxn 1).
- 3) RH elimination (rxn 2) also occurs (when $\text{R} = \text{CH}_3$ and perhaps also C_2H_5 and phenyl) but does not compete with H_2 elimination
- 4) Decompositions of dialkylsilanes and monoalkylsilanes ($\text{R} > \text{CH}_3$) are accelerated by silylene induced chain reactions (e.g. rxns 3-6, for ethylsilane).
- 5) Free radical chain reactions, in addition to silylene chains, occur in the decompositions of dialkyl and higher alkylsubstituted monosilanes.
- 6) Alkyl substitution for H in monosilanes increases decomposition activation energies by about $4 \pm 1 \text{ Kcal/R group}$ up to R_2SiH_2 (i.e. the order of thermal stabilities in monosilanes is $\text{SiH}_4 < \text{RSiH}_3 < \text{R}_2\text{SiH}_2 \approx \text{R}_3\text{SiH}$).
- 7) Disilanes decompose mainly by 1,2 H shifts (e.g. rxn 7), but 1,1 H_2 eliminations become important at higher temperatures (e.g. rxn 8).
- 8) Decompositions by 1,2 H_2 eliminations (e.g. rxn 9) do not occur.

Alkylsilane Decompositions




CHAIN



Alkyldisilane Decompositions Illustrated for a 1,1-alkyldisilane



1,2 H-shift



1,1 H₂ elimination



does not occur

Area #2) Relative Rate Kinetics of the Trapping Reactions of SiH₂.

Systems studied were silylene reactions with ethylene and butadiene¹⁰, silylene reactions with MeOH and butadiene⁹, silylene reactions with acetylene and silane⁴, and silylene reactions with 1-butene and silane¹¹. Systems under investigation are SiH₂ with 1-pentene and cis-2-butene.


Results of the area #2 relative rate studies were:

1) Silylene trapping by multiple bonded substrates are complex processes involving multiple reaction channels, some of which do not produce observable products. For example silylene trapping by acetylene and silane is non-quantitative and mass balances of the observable products, which are disilane, vinylsilane, ethynylsilane, ethylene, ethynylvinylsilane and ethynldivinyisilane, diverge increasingly from theoretical as the reaction proceeds. The mechanism, which involves silacyclopropane formations and subsequent isomerizations and decompositions, is shown in Scheme 1 below. [Observed products are underlined.]




SEARCHED	INDEXED
SERIALIZED	FILED
OCT 1961	
FBI - NEW YORK	
A-1	

Scheme I: Mechanism of the Silylene + Acetylene Reaction



2) Trapping by ethylene and other olefins is similarly complex. The corresponding mechanism shown below in Scheme II:






Scheme II: Mechanism of the Silylene + Ethylene Reaction



The intermediate silylenes of the above reaction (SiH_2 , $\text{C}_2\text{H}_5\text{SiH}$ and $(\text{C}_2\text{H}_5)_2\text{Si}$) can be trapped by Si-H insertion trapping agents to eventually produce alkylated silanes. Also, on the basis of poor product mass balances, it is clear that continued trapping by ethylene leads not only to vinylsilane products but more importantly to polymeric materials. Solid deposits on the reaction cell walls and connecting inlet and outlet tubes are further evidence for polymer formation.

3) Relative rate constant determinations for silylene trapping reactions by various substrates have been determined, and the relevant reactions and their kinetics are shown below.

Competing Reactions



Relative Kinetics

$$\ln(k_{10}/k_{11}) = 5.9 - 5846 \text{ cal/RT}$$

$$\ln(k_{10}/k_{12}) = 1.5 - 712 \text{ cal/RT}$$

$$k_{13}/k_{14} = 2.1 \text{ (643-689 K)}$$

Area #3) Mechanisms and Kinetics of the Isomerizations and Decompositions of Organosilylenes.

Systems studied were the isomerization and decomposition kinetics of ethylsilylene², n-propylsilylene¹, ethynylsilylene⁶, and most importantly, n-butylsilylene⁷, and sec-butylsilylene¹¹].


Results of the area #3 studies are:

- 1) Alkylsilylenes ($R > \text{Me}$) decompose at SPST and static system reaction temperatures to SiH_2 and olefins.
- 2) The mechanism of decomposition involves silacyclopropane intermediates (or perhaps at SPST temperatures, silacyclobutanes).
- 3) Alkylsilylene decompositions have activation energies less than 30 Kcal.
- 4) The decomposition of sec-butylsilylene occurs with an activation energy significantly lower than its decomposition reaction enthalpy. This requires a very loose transition state for the decomposition reaction and also requires that the decomposition occur by a consecutive step process (e.g. a biradical mechanism).
- 5) Silylenes add very readily to pi bonds to form very labile silacyclopropanes. These addition reactions are high A-factor processes, probably consecutive step biradical processes, and cannot be, as formally believed, single step concerted processes.
- 6) Silacyclopropanes have three important and competing reaction pathways: decomposition to silylene and olefin, isomerization to a silylene via an H atom shift from silicon to carbon and

isomerization to a vinylsilane via an H atom shift from carbon to silicon.

7) Alkylsilylenes readily isomerize via intramolecular C-H insertion reactions. The 3-center insertions producing silacyclopropanes are the most facile and are the dominant reactions at 'low' temperatures. Higher center insertions, (e.g. four center to produce silacyclobutanes, and five center to produce silacyclopentanes) appear to occur only under 'high' temperature shock tube reaction conditions. Thus in the reactions of n-butylsilylene generated at shock tube temperatures (i.e. $T \approx 1200$ K), see Scheme III, the rate constant ratios for 3-center/4-center/5-center reactions were calculated on the basis of product yields to be $k_{3\text{-center}}/k_{4\text{-center}}/k_{5\text{-center}} = 1/0.3/1$.⁷

Scheme III: n-Butylsilylene Isomerization and Decomposition Rxns











2. PROGRESS TOWARD UNDERSTANDING THE DETAILS OF THE MECHANISM FOR CARBOSILANE FORMATION FROM ALKYSILANE PYROLYSES.

From the above studies, we have managed to learn enough about the general reaction and kinetic behaviors of the important intermediates and reactive products produced in organosilicon pyrolysis reaction systems to be able to propose a general reaction mechanism for carbosilane production.¹² Thus a mechanism, shown in part in Scheme IV, has been presented relative to the dimethylsilane pyrolysis system. This mechanism explains both qualitatively and semiquantitative the complex of products formed in that system (including 4-membered, 6-membered and 6-membered polycyclic adamantane type carbosilane products). Thus modeling

predictions with regard to the nature and yield of products are in good agreement with the experimental data.

Scheme IV: Mechanism of the Dimethylsilane Pyrolysis



3. OTHER RECENTLY COMPLETED RESEARCH AND OTHER ONGOING RESEARCH

There are several projects still in progress which have been supported by AFOSR. These projects fall under the three research areas described previously and will produce results that are quite significant to those areas. Among these are the kinetic studies on 1,1,2-trimethyldisilane¹⁴, 1,1-dimethyldisilane¹⁵ and 1,1,2,2-tetramethyldisilane.¹³ The data from these studies will be used to resolve long standing problems concerning the heats of formation of alkylated disilanes, and even more important, heats of formation of alkylated silylenes.¹⁶ Preliminary calculations for the latter give $\Delta H_f^\circ(\text{MeSiH}) \approx 48$ Kcal/mol and $\Delta H_f^\circ(\text{Me}_2\text{Si}) \approx 34$ Kcal/mol. In addition, the heat of formation of hexamethyldisilane is calculated to be about -78 Kcal/mol, a value in agreement (within the experimental errors) of the -81 Kcal/mol value deduced from the decomposition kinetics of hexamethyldisilane as reported by Davidson et.al. This work¹⁶, when coupled with other available and reliable literature data, will also result in a

revised additivity scheme for the accurate estimation of the thermodynamic properties of C,H,and Si containing compounds.¹⁷

Work continues on studies of the intramolecular rearrangements and decompositions of long chain alkylsilylenes, where the latter are generated via silylene additions to long chain olefins. The results of these studies will supplement our understanding of alkylsilylene reactions and hopefully clarify the kinetics of their elementary reactions (i.e. their intramolecular insertions, ring openings and decompositions). Thus, for example, while reasonable estimates of the rate constants for intramolecular 3,4,5 and 6 membered ring formations can be made (i.e. such estimates have been used successfully in the various modeling studies mentioned in this report) no direct and unambiguous data have yet been gathered for these reactions. This has been, and will continue to be, one of the main focuses of our ongoing research with or without AFOSR support. A complete knowledge of carbosilane formation kinetics will not be possible until the kinetics of these kinds of reactions are fully understood.

PUBLICATIONS SUPPORTED BY THIS GRANT

1. B.A. Sawrey, H.E. O'Neal, M.A. Ring and D. Coffey Jr., *Int. J. Chem. Kinet.*, 16, 801 (1984).
2. S.F. Rickborn, M.A. Ring and H.E. O'Neal, *Int. J. Chem. Kinet.*, 16, 1371 (1984).
3. R.T. White, R.L. Espino-Rios, D.S. Rogers, M.A. Ring and H.E. O'Neal, *Int. J. Chem. Kinet.*, 17, 1029 (1985).
4. J.W. Erwin, M.A. Ring and H.E. O'Neal, *Int. J. Chem. Kinet.*, 17, 1067 (1985).
5. S.F. Rickborn, D.S. Rogers, M.A. Ring and H.E. O'Neal, *J. Phys. Chem.*, 90, 408 (1986).
6. D.S. Rogers, M.A. Ring and H.E. O'Neal, *Organometallics*, 5, 1521 (1986).
7. B.A. Sawrey, H.E. O'Neal and M.A. Ring, *Organometallics*, 6, 720, (1987).
8. J.G. Martin, M.A. Ring and H.E. O'Neal, *Organometallics*, 5, 1228 (1986).
9. D.S. Rogers, H.E. O'Neal and M.A. Ring, *Organometallics*, 5, 1467 (1986).
10. D.S. Rogers, K.L. Walker, M.A. Ring and H.E. O'Neal, *Organometallics*, 6, 2313 (1987).
11. A.P. Dickenson, K.E. Nares, M.A. Ring and H. E. O'Neal, *Organometallics*, 6, 2596 (1987).
12. H.E. O'Neal and M.A. Ring, *Organometallics*, 7, 0000 (1988), April.
13. K.A. Nares, M.A. Ring, H.E. O'Neal, in preparation.
14. K.A. Nares, M.A. Ring, H.E. O'Neal, to be published.
15. M. Harris, H.E. O'Neal and M.A. Ring, to be published.
16. H.E. O'Neal and M.A. Ring, in preparation.
17. G. Licciardi, H.E. O'Neal and M.A. Ring, to be published.

END

DATE

FILMED

DTIC

6-88